

NIH RELAIS Document Delivery

NIH-10098365

NIH -- W1 CL696

PAMELA GEHRON ROBEY
CSDB/NIDR/NIH Bldng 30 Rm 228
30 CONVENT DRIVE MSC 4320
BETHESDA, MD 20892

ATTN:	SUBMITTED:	2001-12-27 14:47:23
PHONE: 301-496-4563	PRINTED:	2001-12-28 10:20:47
FAX: 301-402-0824	REQUEST NO.:	NIH-10098365
E-MAIL:	SENT VIA:	LOAN DOC 5382029

NIH	Fiche to Paper	Journal
TITLE:	CLINICAL ENDOCRINOLOGY	
PUBLISHER/PLACE:	Blackwell Scientific Publications.	Oxford,
VOLUME/ISSUE/PAGES:	1992 Jun;36(6):605-8	605-8
DATE:	1992	
AUTHOR OF ARTICLE:	Premawardhana LD; Vora JP; Mills R; Scanlon MF	
TITLE OF ARTICLE:	Acromegaly and its treatment in the McCune-Albright	
ISSN:	0300-0664	
OTHER NOS/LETTERS:	Library reports holding volume or year	
	0346653	
	1424186	
SOURCE:	PubMed	
CALL NUMBER:	W1 CL696	
REQUESTER INFO:	AB424	
DELIVERY:	E-mail: probey@DIR.NIDCR.NIH.GOV	
REPLY:	Mail:	

NOTICE: THIS MATERIAL MAY BE PROTECTED BY COPYRIGHT LAW (TITLE 17, U.S. CODE)

-----National-Institutes-of-Health,-Bethesda,-MD-----

Case of the Month

Acromegaly and its treatment in the McCune–Albright syndrome

L. D. K. E. Premawardhana, J. P. Vora, R. Mills* and M. F. Scanlon

Departments of Medicine and *Surgery, University of Wales College of Medicine, Cardiff, UK

(Received 19 August 1991; returned for revision 19 September 1991; finally revised 10 December 1991; accepted 2 January 1991)

Summary

The McCune–Albright syndrome, comprising polyostotic fibrous dysplasia, cutaneous pigmentation and endocrine hyperfunction, is occasionally complicated by acromegaly due to a pituitary adenoma. We report a patient with the McCune–Albright syndrome and acromegaly, who developed secondary hypothyroidism and hypoadrenalism, in whom surgical removal of the pituitary tumour was technically difficult. A combination of a long-acting somatostatin analogue ('Sandostatin') and external irradiation were therefore used as treatment.

Over 50 years ago, McCune and Albright described a syndrome characterized by polyostotic fibrous dysplasia, cutaneous pigmentation and endocrine hyperfunction (McCune, 1936; Albright *et al.*, 1937). Sexual precocity, occurring more frequently in girls, is by far the most common endocrinopathy (Maurag & Blizzard, 1986). Since then, other endocrinopathies have been described in association with the syndrome, including hyperthyroidism, Cushing's syndrome, hyperprolactinaemia, hypophosphataemic rickets and acromegaly. We report a patient with the characteristic triad of the McCune–Albright syndrome who developed acromegaly due to a radiologically confirmed pituitary adenoma.

Case presentation

The patient, a 26-year-old Caucasian female, had left sided nasal obstruction and rhinorrhoea following a left partial maxillectomy under general anaesthesia for the McCune–Albright syndrome which was diagnosed following the onset of intermittent vaginal bleeding and breast enlargement at the age of 3 years. Patchy cutaneous pigmentation and

fibrous dysplasia were observed in early childhood. Menarche was fully established at 8–9 years. During adolescence, shortening of the left femur due to dysplastic changes was treated by multiple osteotomies with bone grafting.

Enquiry revealed enlargement of feet (shoe size (UK) increasing by 2½) and increased sweating for 1 year approximately. There was no headache, visual disturbance, weight gain or galactorrhoea. Menses were regular. There was no relevant family history. The patient was married but had no children.

Examination revealed facial asymmetry with prominence of her left maxilla and mandible. Her skin was greasy and thickened. Areas of flat, irregular cutaneous pigmentation were present bilaterally but did not cross the midline (Fig. 1). However, other features of acromegaly such as acral enlargement, prognathism or prominence of supraorbital ridges, goitre or galactorrhoea were not observed. Visual fields and fundi were normal.

The results of endocrine evaluation were consistent with acromegaly with growth hormone (GH) levels during a day profile ranging between 16.0 and 37.7 mU/l. Glucose tolerance was normal (fasting and 2-hour values being 4.9 and 6.5 mmol/l respectively), but GH levels failed to suppress adequately from a basal value of 34.5 mU/l to a minimum of 14.8 mU/l. TRH produced a paradoxical rise in GH from 27.0 to over 100 mU/l. Basal prolactin (PRL) was 310 mU/l with a normal response to TRH. During insulin-induced hypoglycaemia (plasma glucose 1.8 mmol/l) GH rose from 35 to 52 mU/l and prolactin levels remained stable at approximately 400 mU/l. However, plasma cortisol levels were unrecordable (< 28 nmol/l) during the basal period and following administration of insulin. Other investigations revealed free T₄ 9 pmol/l; TSH 1.5 mU/l; LH 5.4 U/l; FSH 3.5 U/l; 17- β -oestradiol 80 pmol/l and progesterone < 2 nmol/l.

High resolution CT scan of the pituitary fossa with contrast enhancement demonstrated an enhancing pituitary tumour with suprasellar extension with thickened surrounding bone and diffuse vault sclerosis clearly affecting occipital and parietal bones (Fig. 2).

Having confirmed the diagnosis of acromegaly due to a pituitary adenoma, she was started on hydrocortisone replacement therapy. Thereafter a transethmoidal hypophysectomy was undertaken. At operation the enlarged pituitary was surrounded by a 'venous lake' and surgical access was not possible due to haemorrhage. Surgery was therefore abandoned.

Correspondence: Dr L. D. K. E. Premawardhana, Division of Endocrinology, Department of Medicine (7th Floor), University of Wales College of Medicine, Heath Park, Cardiff CF4 4XN, UK.




Fig. 1 Cutaneous pigmentation, with irregular margins in McCune-Albright syndrome. Shortening of the left femur is also visible.

Following attempted surgery, elevated GH levels were confirmed during a day profile and pituitary irradiation was undertaken by means of three-field external deep irradiation therapy. In the interim, GH hypersecretion was treated medically. A single dose of bromocriptine 2.5 mg orally, reduced growth hormone levels from 54.4 to 26.3 mU/l at 2 hours rising to 39.6 mU/l at 4 hours post administration.

By contrast, 100 µg Sandostatin (octreotide, Sandoz) subcutaneously reduced GH levels from 58.8 to 7.1 mU/l at 2 hours and thereafter to 21.0 mU/l at 4 hours post injection.

Persistent secondary hypothyroidism (free T₄ 10 pmol/l and TSH 1.2 mU/l) was thereafter treated with L-thyroxine 100 µg/day. She was commenced on treatment with octreotide 100 µg three times daily subcutaneously and hydrocortisone 10 mg twice daily.

An initial short lived watery diarrhoea settled and she has been free of side-effects. Clinical improvement was noticed by lack of excessive sweating and decrease in the size of her feet.

One year into treatment her GH day profiles showed partial suppression (9.8–13.6 mU/l) but remain high off treatment for a week (32–64 mU/l). CT scans did not show significant change in size of tumour or suprasellar extension. Her liver function tests, thyroid function tests, fasting and 2-hour postprandial blood sugars and HBA1C were completely normal. She was free of symptoms of gall bladder calculi. Her dose of octreotide has now been increased to 150 µg three times daily.

Discussion

Acromegaly is a rare endocrinopathy associated with the McCune-Albright syndrome. Reviews of the literature by Lipson and Hsu (1981) and Cuttler *et al.* (1989) revealed six and 17 cases respectively of well documented clinical and biochemical acromegaly. Our patient fulfilled the biochemical criteria of acromegaly: high daytime GH levels, non-suppression during oral glucose tolerance testing, and increase after TRH. The surprising endocrine findings in our patient were of marked hypoadrenalism (despite successfully undergoing a general anaesthetic prior to presentation), together with subsequent development of hypothyroidism. As far as we are aware this has not been reported in the McCune-Albright syndrome. GH hypersecretion in the McCune-Albright syndrome differs in several respects from that observed in classical acromegaly. Presentation of gigantism/acromegaly is considerably earlier than in classical acromegaly. In one review, 14 out of 17 patients were below 30 years of age (Lipson & Hsu, 1981). Less than half the patients with McCune-Albright syndrome and acromegaly show radiological evidence of a pituitary adenoma compared with 80–90% in classical acromegaly. Of Cuttler's series only three showed suprasellar extension as in our patient. Hyperprolactinaemia was not observed in our patient in contrast to 85% of patients with McCune-Albright syndrome combined with GH hypersecretion. By comparison, hyperprolactinaemia is a feature of 30–40% of patients with classical acromegaly.




Fig. 2 High resolution contrast enhanced pituitary scan showing fibrous dysplasia of the left temporo-parietal-occipital bones, and pituitary tumour with suprasellar extension.

Pituitary surgery in this syndrome is beset with problems mainly related to fibrous dysplasia and the resultant thickening of bones. At surgery our patient proved to have abnormally thickened dysplastic maxillary and ethmoid bones. However, increased vascularity surrounding the pituitary gland prevented surgical intervention.

The ineffectiveness of bromocriptine in completely suppressing GH levels prompted the use of the long acting somatostatin analogue, octreotide (Sandostatin, Sandoz). Geffner *et al.* (1987) first reported the use of somatostatin in acromegaly associated with the McCune-Albright syndrome. Octreotide has a half-life of 110 minute after subcutaneous injection, and an overall duration of effect of about 6–8 hours (Barnard *et al.*, 1986). A single subcutaneous injection of octreotide produces a reduction of GH levels in 30–60 minutes and maximally suppresses GH levels between 2 and 4 hours. Levels may rise to preinjection levels 8 hours after injection (Frohman, 1991). Our patient's data

are consistent with these findings. Although long-term suppressiveness can be predicted by the acute suppressive effects of a single injection—a reduction of 88% at 2 hours in our patient (Lamberts *et al.*, 1989)—the lack of complete suppression after a year of treatment may be due to an inadequate dose or reduced numbers of somatostatin receptors in the GH secreting adenoma of our patient. Ten to 30% of GH secreting tumours are known to have reduced somatostatin receptors (Reubi & Landolt, 1989). The effects of octreotide on tumour shrinkage are less predictable. Although both microadenomas and macroadenomas have been known to shrink, only about half of all GH secreting tumours decrease in size (Frohman, 1991). Our patient's tumour has not changed in size. We used radiotherapy in addition to octreotide to prevent tumour expansion, particularly in view of the suprasellar extension.

In conclusion, we suggest that the combination of a somatostatin analogue and radiotherapy may be the treat-

ment of choice in patients with McCune-Albright syndrome and acromegaly, in view of the technical difficulties associated with pituitary surgery in this syndrome.

References

- Albright, F., Butler, A.M., Hampton, A.O. & Smith, P. (1937) Syndrome characterized by osteitis fibrosa disseminata, areas of pigmentation and endocrine dysfunction, with precocious puberty in females. *New England Journal of Medicine*, **216**, 727-746.
- Barnard, L.B., Grantham, W.G., Lamberton, P., O'Dorioso, T.M. & Jackson, I.M.D. (1986) Treatment of resistant acromegaly with a long acting somatostatin analogue (SMS 201-995). *Annals of Internal Medicine*, **105**, 856-861.
- Cuttler, L., Jackson, J.A., Uz-Zafar, M.S., Levitsky, L.L., Mellinger, R.C. & Frohman, L.A. (1989) Hypersecretion of growth hormone and prolactin in McCune-Albright syndrome. *Journal of Clinical Endocrinology and Metabolism*, **68**, 1148-1154.
- Frohman, L.A. (1991) Therapeutic options in acromegaly. *Journal of Clinical Endocrinology and Metabolism*, **72**, 1175-1181.
- Geffner, M.E., Nagel, R.A., Dietrich, R.B. & Kaplan, S.A. (1987) Treatment of acromegaly with a somatostatin analogue in a patient with McCune-Albright syndrome. *Journal of Paediatrics*, **111**, 740-743.
- Lamberts, S.W.J., Van Koetsveld, P. & Hofland, L. (1989) A close correlation between the inhibitory effects of insulin-like growth factor I and SMS 201-995 on growth hormone release by acromegalic pituitary tumours *in vitro* and *in vivo*. *Clinical Endocrinology*, **31**, 401-410.
- Lipson, A. & Hsu, T. (1981) McCune-Albright syndrome associated with acromegaly: report of a case and review of literature. *Johns Hopkins Medical Journal*, **149**, 10-14.
- Maurag, N. & Blizzard, M. (1986) The McCune-Albright syndrome. *Acta Endocrinologica*, **113** (Suppl.), 207-217.
- McCune, D.J. (1936) Osteitis fibrosa cystica. The case of a 9-year-old girl who also exhibits precocious puberty, multiple pigmentation of the skin and hyperthyroidism. *American Journal of Diseases of Children*, **52**, 743-744.
- Reubi, J.C. & Landolt, A.M. (1989) The growth hormone responses to octreotide in acromegaly correlate with adenoma somatostatin receptor status. *Journal of Clinical Endocrinology and Metabolism*, **68**, 844-850.

Commentary

Giovanni Faglia

Institute of Endocrine Sciences, University of Milan, Italy

The Case of the Month (Premawardhana et al., 1992) reports on a patient with the McCune-Albright syndrome in whom endocrine hyperfunction was expressed as precocious puberty and acromegaly. Unusually, secondary hypoadrenocorticism and hypothyroidism were present probably due to pituitary insufficiency caused by a huge GH-secreting pituitary adenoma. As the patient could not be operated on, she was treated with octreotide (which caused serum GH to fall by about 80%) in combination with radiotherapy.

Recently, Weinstein et al. (1991) have demonstrated the presence of somatic activating mutations in the α -chain of the stimulatory G protein—identical to those previously shown in some functioning endocrine tumours (Landis et al., 1989; Lyons et al., 1990; Suarez et al., 1991)—in several tissues of four patients with McCune-Albright syndrome. These mutations cause substitution of other aminoacids for Arg 201 of $G_s\alpha$ resulting in the constitutive activation of adenylyl cyclase. As all the diverse syndromes of endocrine hyperfunction described in the McCune-Albright syndrome involve cells that respond to stimulatory hormones via the activation of adenylyl cyclase system and the subsequent cAMP formation as second messenger, it is likely that these mutations are responsible for unstimulated hormone hypersecretion (Levine et al., 1991).

The subset of acromegalic patients with mutated $G_s\alpha$ pituitary GH-secreting tumours respond particularly well to octreotide treatment (Spada et al., 1990). In view of this, it

is reasonable to assume that octreotide is the first choice therapy for acromegaly associated with the McCune-Albright syndrome, and it is tempting to speculate that this drug might also control other associated endocrine hyperfunctions, provided that the involved glands possess somatostatin receptors.

References

- Landis, C.A., Masters, S.B., Spada, A., Pace, A.M., Bourne, H.R. & Vallar, L. (1989) GTPase inhibiting mutations activate the α chain of G_s and stimulate adenylyl cyclase in human pituitary tumours. *Nature*, **340**, 692-696.
- Levine, M.A. (1991) The McCune-Albright syndrome. The whys and wherefores of abnormal signal transduction. *New England Journal of Medicine*, **325**, 1738-1740.
- Lyons, J., Landis, C.A., Harsh, G., Vallar, L., Grunewald, K., Feichtinger, H., Duh, Q.Y., Clark, O.M., Kawasaki, E., Bourne, H.R. & McCormick, F. (1990) Two G protein oncogenes in human endocrine tumors. *Science*, **249**, 655-659.
- Premawardhana, L.D.K.E., Vora, J.P., Mills, R. & Scanlon, M.F. (1992) Acromegaly and its treatment in the McCune-Albright syndrome. *Clinical Endocrinology*, **36**, 605-608.
- Spada, A., Arosio, M., Bochicchio, D., Bazzoni, N., Vallar, L., Bassetti, M. & Faglia, G. (1990) Clinical, biochemical and morphological correlates in patients bearing growth hormone-secreting pituitary tumours with or without constitutively active adenylyl cyclase. *Journal of Clinical Endocrinology and Metabolism*, **71**, 1421-1426.
- Suarez, H.G., du Villar, J.A., Caillou, B., Schlumberger, M., Parmentier, C. & Monier, R. (1991) *gsp* mutations in human thyroid tumours. *Oncogene*, **6**, 677-679.
- Weinstein, L.S., Shenker, A., Gejman, P.V., Merino, M.J., Friedman, E. & Spiegel, A.M. (1991) Activating mutations of the stimulatory G protein in the McCune-Albright syndrome. *New England Journal of Medicine*, **325**, 1688-1695.